

THE EFFECT OF RATION WITH ANTIBIOTICS (Virginiamycin) and TEMULAWAK (Curcuma xanthorriza Roxb.) TO BROILER PERFORMANCES

Lovita Adriani¹, Endang Sujana², Andi Mushawwir³

^{1,3} Laboratorium Fisiologi Ternak dan Biokimia Fakultas Peternakan Universitas Padjadjaran
Kampus Jatinangor Sumedang Jabar Indoensia.

Email : andi_mh@unpad.ac.id

² Laboratorium Produksi Ternak Unggas Fakultas Peternakan Universitas Padjadjaran Kampus
Jatinangor

ABSTRACT

The research about “The Effect of Ration with Antibiotics (Virginiamycin) and Temulawak (Curcuma xanthorriza roxb.) to Broiler Performance ” has been conducted at Faculty of Animal Husbandry, Padjadjaran University, Jatinangor, that started from 11st of March to 15th of April, 2009. The purpose of this research is to learn about the effect of Virginiamycin and Temulawak addition to performance . The research used 100 broiler chicken from Cobb CP 707 strain. The research used Completely Randomized Design (RAL-Rancangan Acak Lengkap) with four treatments. R0: Control,; R1: ration with 15 ppm Virginiamycin; R2: ration with 9% Temulawak; R3: ration with 15 ppm Virginiamycin and 9% Temulawak. Duncan’s Multiple Range Test is used to indicate the difference between treatments. Variable being observed is performance, and income over feed cost. Result of the research showed that temulawak and virginiamycin addition does not provide significant difference on performance feed consumption, weight gain,

Keywords: *Virginiamycin, temulawak, weight gain, carcass final weight , broilers*

INTRODUCTION

The preventive diseases and enhancement of growth, feed intake and feed efficiency are importance factor. Most antibiotic as a feed additive requires alternative ways to stabilize the health and growth performance as antibiotic as a feed additive Virginiamycin as antibiotic used to growth promoter. Antibiotic is feed additive

Used virginiamycin in true composition, effected to restraining pathogen bacteria population in intestinal and decreasing negative efect of normal flora population that over in the gastrointestinal.

Combinations of vitamin and amino acid which is be able do together to prevent micro nutrient deficiency, increasing growing and not caused if used at all times. Despitefully of antibiotic, utilizing herbal be also a choice as feed additive.

Utilizing herbal contribute to increasing appetite and also growing. Herbal used occasionally is *curcuma xanthorhiza* (temulawak). Temulawak as far back as medicine concoction matter. Contents of temulawak consist of *curcumin*, colagoga, atsiri oil (volatile oil).

Curcumin is substance that protected of oxidasi on eritrocite and haemoglobin by nitrite compounds. *Curcumin* can be increasing hepathoglobin and haemopexin protein synthesis in the liver, so lead which is bonding with haemoglobin be able to destruction in the liver. Kolagoga is substance that to increasing role of gall and decreasing body fat, so we get low fat meat and high meat composition.

Volatile oil also called essentia oils, that formed at reticulum endoplasmic in the plan cells, and providable by distillation. Volatile oil not effected on microorganism population, but positive effected on digestion enzymes. Now, volatile oils getting popular in the agriculture and animal livestock sector, because this oils used as digestion and metabolism promoters, and not giving rise to resistance in animal.

Based on description above, so need to do research The Effect of Ration with Antibiotics (Virginiamycin) and Temulawak (*Curcuma xanthorriza roxb.*) to Broiler Performance.

MATERIALS AND METHODS

The experiment was carried out in the Experimental poultry unit of the Faculty of Animal Husbandry, Padjadaran University. One hundred, day old chick, Cobb broiler were chosen. The research used Completely Randomized Design (CRD).. They were consisting

four different treatments, with four replications in one treatment. Chicken were fed ad libitum with four ration. The composition of the ration were :

R₀ = Control

R₁ = Ration with 15 ppm virginiamycin

R₂ = Ration with 9 % temulawak

R₃ = Ration with 15 ppm virginiamycin and 9% temulawak

The analysis of the treatment were body weight, and carcass weight,

RESULT AND DISCUSSION

1. The Effect of Treatment on Final Body Weight in Broiler

Tabel 1. Average final body weight on each treatment

Replication	Treatment			
	R0	R1	R2	R3
(g).....			
1	1320	1300	1180	1300
2	1320	1280	1270	1120
3	1500	1300	1420	1300
4	1160	1320	1580	1400
5	1200	1250	1250	1540
Total	6500	6450	6700	6660
Average	1300	1290	1340	1332

In Table 1. are presented the results of the treatment, the data were followed with statistic analysis. Results indicated that all treatments have no significantcy on the final body weight. 9% **Curcuma xanthorrhiza Roxb/** temulawak) combined with 15 ppm virginiamycin, have no significancy, even the. Virginiamycin effectively as anti bacteria and content cystein and lysine (Komisi obat Hewan, Departemen Pertanian, 2006). According to Rofiq, 2003, using 15 ppp virginiamycin will affected the gastrointestinal tract, and the ration added with antibiotic for a long term even in low dosage will disturb the balance of acid-base in small intestine and

intestinal damage, also affected the nutrient absorption at least will result in the lower of final body weight. .. According to Lesson and Summers, 2001), virginiamycin will change the intestinal microflora, and the ceca will be oedema and filled with humid excrete and the epithelium will lysis.

Antimicrobial growth promoters (virginiamycin) are antibiotics added to the feed of food animals to enhance their growth rate and production performance. The mechanism by which virginiamycin work is not clear. Virginiamycin reduce normal intestinal flora and harmful gut bacteria. The effect on growth may be due to a combination of both fewer normal intestinal flora and fewer harmful bacteria.

The combination between virginiamycin and temulawak has advantages, because the anti-inflammatory effect, reduce the diarrhea effect and to improve the villi and mucosa structure, and the inflammation of intestine.

2 . The influence of the treatment on the broiler carcass weight

Tabel 2.. The influence of the treatment on the broiler carcass weight

Replication	Treatment			
	R0	R1	R2	R3
(g).....			
1	850	900	700	900
2	900	770	700	650
3	1000	750	825	870
4	700	700	1000	900
5	750	700	1200	1000
Total	4200	3820	4425	4320
Average	840	764	885	864

In Table 2.. results indicated that 9% virginiamycin in ration (R1), have the lowest average than other treatment (R2 and R3) although have no significant on the final body weight. The function of colagoga in temulawak in ration (R2), is to change the lipid became energy and increased the muscle, at least could increased the carcass weight.

The choloretic properties and cinarina colagoga, could stimulates the production of bile in the liver and facilitates the clearing later in the gallbladder, which helps digestion of fats. Bile, made up of bile salts and cholesterol, is secreted by hepatocytes and is stored in the gallbladder. It is excreted after ingestion of food to metabolize and digest. By action of bile, fats from oily foods and fried foods are emulsified (broken up into small molecules), transformed into droplets that are degraded by pancreatic and intestinal lipases, being apt to be degraded by enzymes secreted by lipases pancreas. The bile, prepared digestion of fatty substances and later digestion occurs through the pancreatic juice, the only way to proceed with the breakdown of fats. Digestive enzymes are secreted by the pancreas and comprise lipases, colesterolasas, glucidasas and proteases. (Doymafarma, 2009). The use of choloretic and Bile means in case of non-ulcer dyspepsia and there are heavy and slow digestion

CONCLUSION

Temulawak and virginiamycin has positive response to body weight, and carcass weight. R2 (9% temulawak) and R3 (9% temulawak and 15pp virginiamycin), was better than R0 (control) and R1 (virginiamycin 15ppm)

REFERENCES

- Departemen Kesehatan, 1987. *Formularium Obat Tradisional Indonesia*, Departemen Kesehatan Republik Indonesia, Jakarta.
- Departemen Pertanian, 2006. *Peggunaan Pemacu Pertumbuhan pada Ternak*. . www.wikipedia.com (di akses 29 juli 2008).
- Henderawati A. 1999. *Penurunan Kadar Kolesterol Daging Broiler Dengan Penambahan Temulawak (Curcuma Xanthorrhiza Roxb.) Dalam ransum*. Thesis. Fakultas Pasca Sarjana IPB. Bogor
- Infonet. 1999. Antibiotik; Dulu Antibiotik, Kini Probiotik. Majalah edisi 067. Asahi. Jakarta.
- http://www.doymafarma.com/daymafarma/ctl_Servlet?_f=37&ld=13053415

1. Leclercq R, Derlot E, Weber M, Duval J, Courvalin P. Transferable vancomycin and teicoplanin resistance in *Enterococcus faecium*. *Antimicrob Agents Chemother* 1989;33:10-5.
2. Noble WC, Virani Z, Cree RG. Co-transfer of vancomycin and other resistance genes from *Enterococcus faecalis* NCTC 12201 to *Staphylococcus aureus*. *FEMS Microbiol Lett* 1992;72:195-8.
3. Berchieri A. Intestinal colonization of a human subject by vancomycin-resistant *Enterococcus faecium*. *Clin Microbiol Infect* 1999;5:97-100.
4. Jensen LB, Ahrens P, Dons L, Jones RN, Hammerum A, Aarestrup FM. Molecular analysis of the Tn1546 from vancomycin resistant enterococci isolated from animals and humans. *J Clin Microbiol* 1998;36:437-42.
5. Jensen LB. Differences in the occurrence of two base pair variants of Tn1546 from vancomycin-resistant enterococci from humans, pigs and poultry. *Antimicrob Agents Chemother* 1998;42:2463-4.
6. Thal L, Donabedian S, Robinson-Dunn B, Chow JW, Dembry L, Clewell DB, et al. Molecular analysis of glycopeptide-resistant *Enterococcus faecium* isolates collected from Michigan hospitals over a 6-year period. *J Clin Microbiol* 1998;36:3303-8.
7. Bager F, Aarestrup FM, Madsen M, Wegener HC. Glycopeptide resistance in *Enterococcus faecium* from broilers following discontinued use of avoparcin. *Microb Drug Resist* 1999;5:(in press).
8. Klare I, Badst, bner D, Konstabel C, B.hme G, Claus H, Witte W. Decreased incidence of VanA-type vancomycin-resistant enterococci isolated from poultry meat and from fecal samples of humans in the community after discontinuation of avoparcin usage in animal husbandry. *Microb Drug Resist* 1999;5 (in press).
9. Welton LA, Thal LA, Perri MB, Donabedien S, McMahon J, Chow JW, et al. Antimicrobial resistance in enterococci isolated from turkey flocks fed virginiamycin. *Antimicrob Agents Chemother* 1998;42:705-8.
10. Van den Bogaard AE, Mertens P, London NH, Stobberingh EE. High prevalence of vancomycin- and pristinamycin-resistant enterococci in healthy humans and animals in The Netherlands: is the addition of antibiotics to animal feed to blame? *Antimicrob Agents Chemother* 1997;40:454-6.
11. Aarestrup FM, Bager F, Madsen M, Jensen NE, Meyling A, Wegener HC. Surveillance of antimicrobial resistance in bacteria isolated from food animals to growth promoters and related therapeutic agents in Denmark. *APMIS* 1998;106:606-22.
12. Hammerum AH, Jensen LB, Aarestrup FM. Detection of the SatA gene and transferability of virginiamycin resistance in *Enterococcus faecium* from food animals. *FEMS Microbiol Lett* 1998;168:145-51.
13. Jakobsen BM, Skou M, Hammerum AM, Jensen LB. In vivo transfer of the satA gene between isogenic strains of *Enterococcus faecium* in the mammalian gastrointestinal