

UNIVERSITATEA DE ȘTIINȚE AGRICOLE ȘI MEDICINĂ VETERINARĂ
ION IONESCU DE LA BRAD
IAȘI-ROMANIA

LUCRĂRI ȘTIINȚIFICE

SERIA MEDICINĂ VETERINARĂ
VOL. 56 (nr. 3-4)



Editura „ION IONESCU DE LA BRAD“
ISSN 1454-7406

**UNIVERSITATEA DE ȘTIINȚE AGRICOLE ȘI MEDICINĂ
VETERINARĂ "ION IONESCU DE LA BRAD"
IAȘI – ROMÂNIA**

LUCRĂRI ȘTIINȚIFICE

SERIA MEDICINĂ VETERINARĂ

VOL. 56 (3 – 4)

ISSN 1454-7406

**Editura „ION IONESCU DE LA BRAD“ Iași
2013**

COLEGIUL DE REDACȚIE / EDITORIAL BOARD

Redactor șef / Editor in Chief - Mihai Mareș

Secretar de redacție / Secretary - Valentin Năstasă

Membru / Member - Mariana Grecu

COMISIA DE REFERENȚI / ADVISORY BOARD

Prof. dr. Liviu Miron – USAMV Iași (Romania)

Prof. dr. Gheorghe Solcan – USAMV Iași (Romania)

Prof. dr. Gheorghe Savuța – USAMV Iași (Romania)

Prof. dr. Gabriel Predoi – USAMV București (Romania)

Prof. dr. Cornel Cătoi – USAMV Cluj-Napoca (Romania)

Prof. dr. Viorel Herman – USAMV Timișoara (Romania)

Assoc. Prof. Dorina Timofte Carter – University of Liverpool (UK)

Assoc. Prof. Valentin Năstasă – USAMV Iași (Romania)

Assoc. Prof. Mihai Mareș - USAMV Iași (Romania)

Responsabilitatea privind conținutul articolelor, inclusiv traducerea acestora în limba engleză, revine exclusiv autorilor.

The entire responsibility for the content of papers, including the English translation, belongs to the authors.

CUPRINS

THE USE OF IOMERON 350 IN CAT UROGRAPHY Radu Lăcătuș, Robert Cristian Purdoi, Ioan Sălăgean, Ionel Papuc	215 - 221
A STUDY ON THE RELATION BETWEEN THE SIZE AND SEX OF DOG, AND THE TSH, THYROXINE AND TRIIODOTHYRONINE SERUM CONCENTRATION IN CANINE POPULATION IN MUSCEL AREA Gabriel I. Vișoiu, Victor Crivineanu, Gheorghe V. Goran	222 - 226
THE INCIDENCE OF TOPOGRAPHIC GASTROINTESTINAL DISORDERS AT PET CARNIVORES V. Boghian	227 - 232
CLINICAL AND PARACLINICAL ASPECTS IN TOPOGRAPHIC GASTROINTESTINAL DISORDERS IN DOG: GASTRIC TORSION, INTUSSUSCEPTION AND INTESTINAL VOLVULUS V. Boghian	233 -237
RABBIT GENERAL ANESTHESIA FOR ENUCLEATION. CASE STUDY Alexandru Cosmin Tutunaru, Florin Leau, Alexandru Șonea, Charlotte Sandersen	238 - 241
CHIARI-LIKE, MENINGOENCEPHALITIS AND ATLANTOAXIAL SUBLUXATION IN A YORKSHIRE TERRIER DOG Florin Eugen Grosu, Panagiotis Mantis	242 - 244
OBSERVATIONS REGARDING VAGINAL PROLAPSE IN BITCH – CASE STUDY Roșca P., Drugociu D., Ciornei Șt.G., Nechifor F., Văleanu (Neculai) Sabina, Ibănescu I.	245 - 249
PROSPECTIVE STUDY REGARDING INCIDENCE OF HIP DYSPLASIA IN FIVE BREEDS IN ROMANIA Aurel Grosu, Ana-Maria Daneliuc, Gentiana Grosu, Florin Eugen Grosu	250 - 252
EFFECTS OF PUERPERAL ENDOMETRITIS ON THE OVARIAN ACTIVITY OF MILK COWS DURING THE POST PARTUM PERIOD Florin Cătălin MLAGIU	253 - 261
IMPORTANCE OF BLOOD INDICATORS OF HEPATIC FUNCTION ON THE REPRODUCTIVE PERFORMANCE AFTER TREATMENT OF CYSTIC OVARIAN DISEASE WITH GONADOTROPHIN-RELEASING HORMONE IN DAIRY COWS Borș S.I., Ruginosu Elena, Creangă Șt., Elena Lopatnicu, Dascălu L.	262 - 266

SUBTOTAL INTRACAPSULAR RESECTION OF PROSTATE ADENOMA WITH TOTAL ABLATION OF THE INTRAPROSTATIC URETHRA IN DOGS Bogdan Alexandru Vițălaru, Ion Dragomirișteanu, Ion Alin Bîrțoiu, Cătălin Pandelaș, Mihaela-Alina Florea	267 - 273
COMPARATIVE EVALUATION ON THE EFFICIENCY AND TOLERABILITY OF THREE NSAIDs USED IN LOCOMOTIVE OSTEOARTICULAR INFLAMMATIONS IN DOG Grecu Mariana, Nastasa Valentin, Musca Raluca, Diana-Luminita Hritcu	274 - 282
VARIATIONS ON CYATHOSTOMINS POPULATION AMONG HORSES FROM DIFFERENT PARTS OF IAȘI AND NEAMȚ COUNTIES C. T. Covasă, L.D. Miron	283 - 289
THE BACTERIAL FLORA OF THE UTERUS ACCORDING TO PARTURITION TYPE IN COWS Florin Nechifor, Dan Drugociu, Petru Roșca, Ștefan Ciornei, Iulian Ibănescu	290 - 297
REMARKS CONCERNING THE PHYSIOLOGICAL CONSTANTS IN COWS DURING PUERPERIUM Florin Nechifor, Dan Drugociu, Petru Roșca, Ștefan Ciornei, Iulian Ibănescu	298 - 302
POST-TRAUMATIC SPINAL HAEMATOMA WITHOUT OSSEOUS LESIONS IN A DOG Adina Zbângu, Mihaela Armasu, M. Musteată, Gh. Solcan	303 - 308
USE OF ACRIDINE ORANGE AND BABES-PAPANICOLAU STAINING FOR THE DIFFERENTIATION OF VARIOUS STAGES OF THE PARASITE <i>BLASTOCYSTIS HOMINIS</i> Doina-Simona Grecu (Mătiut), Maria Larisa Parasca (Ivănescu), Elena-Andreea Hărmănescu, Ioan Moglan, Liviu Miron	309 - 314
THE CHARACTERIZATION OF MIDDLE LATENCY AUDITORY RESPONSES RECORDED WITH SURFACE ELECTRODES IN DOGS WITH DIFFERENT INTRACRANIAL LESIONS Mihaela Armasu, Mihai Musteată, Gabriela Dumitrita Stanciu, Gheorghe Solcan	315 - 319
INFLUENCE OF MICROCLIMATE PARAMETERS IN SWINE REPRODUCTION Loredana Mihaela Vasile, Al. Șonea, Cristinel Șonea, I.Radoi, Catalina Posea	320 - 325
HEPATIC PROFILE OF DAIRY COWS IN RESPONSE TO A TREATMENT OF LIVER DISEASES WITH A PROPYLENE GLYCOL BASED PRODUCT Elena Lopatnicu, Silviu Bors, Gheorghe Solcan	326 - 329
INTERNATIONAL TRAVEL INCREASE AND MALARIA IMPORTATION IN ROMANIA, 2007-2012 Larisa Parasca (Ivănescu), Simona Grecu (Mătiut), Liviu Miron	330 - 339

LOCAL OZONE MONOTHERAPY IN SKIN INFECTION IN A DOG – CASE REPORT B. St Rugină, L. C. Burtan, Ioana Burcoveanu, Cristiana Rugină	340 - 344
THE APPLICABILITY OF SEMEN COLLECTION IN DRONES OF <i>APIS MELIFERA CARPATICA</i> Stefan G. Ciornei, Liviu Runceanu, Dan Drugociu, Petru Roșca, Florin Nechifor	345 - 350
ANALYSIS OF SOME HEMATOLOGICAL AND METABOLIC PARAMETERS IN THE CONDITIONS PROVIDED BY TESTING OF ANTIDEZENTER ON HEALTHY SUCKLING PIGLETS Adrian Vlasiu, Laurenț Ognean, Viorica Chiurciu, Constantin Chiurciu, Florin Zăvoiu, Sebastian Trîncă, Cristina Todoran, Radu Harșan	351 - 358
FAUNE ECTOPARASITAIREET INDICES ÉPIDÉMIOLOGIQUES DU TILAPIA DU NIL <i>OREOCHROMIS NILOTICUS</i> ÉLEVÉ EN ZONE DES LAGUNES (CÔTE D'IVOIRE) Kone Mamadou, Soric Ramona Elena, Cisse Moussa, Affourmou kouassi Frédéric, Ouattara Mamadou, Fantodji Agathe, Miron Dan Liviu	359 - 368
EPIDEMIOLOGICAL STUDY OF SPONDYLOPATHIES IN DOGS C. Daraban, V. Tipișcă, C. Barbazan, A. Băisan, E. Gavrilaş, V. Vulpe	369 – 372
NORMAL ANATOMY OF DOG POPLITEAL LYMPH CENTER, ANATOMICAL VARIANTS AND NON - INVASIVE ASSESSMENT USING ULTRASOUND TECHNIQUES Stan Florin, Damian A., Gudea A., Dezdrobitu C., Delia Bob, Martonoș C., Lăcătuș R., Purdoiu R., Papuc I., Bochiș Ileana	373 - 380
RADIOGRAPHIC DIAGNOSTIC IN TRACHEAL DISORDERS IN SMALL ANIMALS Cristina Barbazan, Andreea Marținaș, A. Baisan, V. Vulpe	381 - 385
A STUDY ON THE PREVALENCE OF DIROFILARIA IMMITIS INFESTATION ON STRAY DOGS IN GALAȚI COUNTY Lavinia Ciucă, Dumitru Acatrinei, Ștefania Merticariu, Liviu Miron	386 - 391
GLYCAEMIC CURVE ASSESSMENT, A MONITORING TOOL FOR ADEQUATE INSULIN THERAPY FOR DIABETES MELLITUS IN CATS Madalina Rosca, Luminita Diana Hritcu, G. Solcan	392 - 398
THE EFFECTS OF CURCUMA (<i>CURCUMA LONGATA</i>) AS NATURAL TENDERIZER ON POULTRY MEAT Hendronoto A.W. Lengkey, Wendry S. Putranto, Eka Wulandari	399 - 402
THE USE OF ACTINOMICETES PRODUCTS IN THE FIGHT AGAINST AMERICAN FOULBROOD Starciuc N., Postolachi Olga, Burțeva Svetlana, Osadci Natalia, Bugneac Veronica, Ciuclea A.	403 - 407

INVESTIGATION OF ANTIMICROBIAL ACTIVITY OF SOME <i>LACTOBACILLUS SALIVARIUS</i> STRAINS ISOLATED FROM DENTAL ROOT CANAL AND OF TWO PROBIOTIC <i>LACTOBACILLUS</i> STRAINS BY INTESTINAL ORIGIN Anca Alexandra Dobrea (Popescu), Constantin Savu, Mimi Dobrea	408 - 411
REPORT CONCERNING RESULTS OF PROFICIENCY TESTING LABORATORY ON ASSAY OF TOBRAMYCINE AND NYSTATIN BY MICROBIOLOGICAL METHOD Simona Sturzu, Daniela Tirsinoaga, Ioana Tihulca, Alina Karina Draghici	412 - 414
PERSPECTIVE EUROPENE IN DOMENIUL PRODUSELOR MEDICINALE VETERINARE Simona Sturzu, Mirela Marinescu	415 - 417
DATA REQUIREMENTS FOR REMOVING THE TARGET ANIMAL BATCH SAFETY TEST (TABST) FOR IMMUNOLOGICAL VETERINARY MEDICINAL PRODUCTS IN THE EU Marius Bunea, Mirela Marinescu	418 - 420
PHYLOGENETIC ANALYSIS OF THE FELINE CORONAVIRUSES FOUND IN HEALTHY CATS AND IN CATS WITH FELINE INFECTIOUS PERITONITIS BASED ON THE NUCLEOCAPSID PROTEIN Ivona Laiu, Lidia Duarte, Cristina Horhoge, Gheorghe Solcan, Aurelian-Sorin Pasca, Mihai Carp-Cărare, Savuta Gheorghe, Sophie LePoder	421 - 426
DETECTION OF THE FELINE CORONAVIRUS IN THE ORGANS OF CATS WITH FELINE INFECTIOUS PERITONITIS USING RT-PCR METHODS Ivona Laiu, Lidia Duarte, Cristina Horhoge, Constantin Pavli, Cătălin Carp-Cărare, Mihai Carp-Cărare, Gheorghe Savuta, Sophie LePoder	427 - 431
METHODOLOGY OF THE ENVIRONMENTAL RISK ASSESSMENT RELATED WITH THE USE OF VETERINARY MEDICINAL PRODUCTS Ioana Valentina Tihulca	432 - 437
APPLICATION OF A REAL-TIME PCR FOR QUANTITATIVE DETECTION OF <i>CAMPYLOBACTER JEJUNI</i> IN FRESH MEAT Vlad-Sabie A., Floriștean V., Borș S.I., Crețu C., Carp-Cărare C., Carp-Cărare M.	438 - 441
EPIDEMIOLOGICAL INVESTIGATIONS ON CLASSICAL SWINE FEVER EVOLUTION IN PIGS AND WILD BOARS IN IAȘI COUNTY Emilia Ion - Popa, Adriana, Anită, Savuța Gheorghe	442 - 446
CLASSICAL SWINE FEVER VIROLOGICAL SURVEILLANCE ON WILD BOARS IN SUCEAVA COUNTY Emilia Ion - Popa, Adriana Anită, Savuța Gheorghe	447 - 450

IMPACT OF FEED CONTAMINATION WITH MYCOTOXINS (OTA AND ZEA) ON PIG AND HUMAN HEALTH Catalina Posea, A. Sonea, A. Bîrtoiu, Monica Roman, Mihaela Vasile	451 - 461
<i>MYCOBACTERIUM AVIUM</i> SSP. <i>AVIUM</i> STUDIED IN NATURALLY INFECTED HENS BY CULTURE AND PCR IDENTIFICATION Macovei Ina Iuliana, Olaru-Péter Simona, Pavel Ionuț, Pașca Aurelian Sorin, Savuța Gheorghe	462 - 467
COMPARATIVE EVALUATION OF THE ANTIFUNGAL ACTIVITY OF VORICONAZOLE AND A NEW PROPICONAZOLE DERIVATIVE (MXP-4509) AGAINST 278 <i>CANDIDA ALBICANS</i> CLINICAL ISOLATES Ramona Moraru, Bogdan Minea, Valentin Năstasă, Bogdan Doroftei, Mihai Mares	468 - 473
COMPARATIVE EVALUATION OF THE ANTIFUNGAL ACTIVITY OF VORICONAZOLE AND A NEW PROPICONAZOLE DERIVATIVE (MXP-4509) AGAINST FLUCONAZOLE-RESISTANT YEAST ISOLATES Ramona Moraru, Bogdan Minea, Valentin Năstasă, Bogdan Doroftei, Mihai Mares	474 - 480
EPIDEMIOLOGICAL INVESTIGATIONS ON ANTIBODIES AGAINST BHV-1 IN FARMS FROM VASLUI COUNTY Anită D., Gramaticu Monica, Anderco Stefania, Anită Adriana	481 - 485
DENTAL MALOCCLUSION IN GUINEA PIG - CASE REPORT Oana Tanase, Constantin Pavli, Florentina Bocaneti	486 - 489
DETECTION OF BOVINE PAPILLOMAVIRUS TYPE 1 IN CUTANEOUS FIBROPAPILLOMAS IN CATTLE Florentina Bocaneti, Maria Angelica Ramos Da Silva, Gennaro Altamura, Annunziata Corteggio, Franco Roperto, Constantin Daraban, Elena Velescu, Giuseppe Borzacchiello	490 - 493
SEROLOGICAL REACTIVITY TO <i>RICKETTSIA CONORII</i> , THE ETIOLOGIC AGENT OF MEDITERRANEAN SPOTTED FEVER IN DOGS FROM ROMANIA Alina Oana Cojocaru (Cas. Paduraru), Gheorghe Savuta	494 - 498
OXIDATIVE STRESS INDUCES THE PRODUCTION OF SPECIFIC MICROPARTICLES AND PROMOTES CELL DYSFUNCTION IN EXOCRINE PANCREAS Andrei Alexandru Constantinescu, Elhassan Yala, Fatiha Zobairi, Florence Toti, Laurence Kessler, Ioan Liviu Mitrea	499 - 509
EXOCRINE MICROPARTICLES PRODUCED IN RESPONSE TO OXIDATIVE STRESS PROMOTE CELL ALTERATION IN ENDOCRINE PANCREAS Andrei Alexandru Constantinescu, Elhassan Yala, Celine Gleizes, Fatiha Zobairi, Florence Toti, Laurence Kessler, Ioan Liviu Mitrea	510 - 519

THE EFFECTS OF CURCUMA (*CURCUMA LONGATA*) AS NATURAL TENDERIZER ON POULTRY MEAT

Hendronoto A.W. Lengkey¹ Wendry S. Putranto¹ and Eka Wulandari¹

¹Universitas Padjadjaran, Bandung, Indonesia

lengkeyhendronoto@gmail.com

Abstract

Forty day old chicks Arbor Acres CP-707 were used randomly in this experiment, to study the effects of various levels of Curcuma (*Curcuma longata*) as natural tenderizer on poultry meat, were studied for six weeks. Research using Completely Randomized Design (CRD). The dietary treatments are: R0 basal diet as control (0% curcuma meal), R1 basal diet + 2% curcuma meal, R2 basal diet + 4% curcuma meal and R3 basal diet + 6% curcuma meal, and each treatment were repeated five times. Results indicated that for the meat fat in broiler which was added curcuma meal in the ration will have less fat to the broiler meat (1.65% - 1.92%) versus 1.99% for basal diet. And for the meat tenderness; the highest meat tenderness was get from the broiler with basal diet + 6% curcuma meal (187,2 mm/g/10sec) and the lowest was get from the broiler with basal diet + 0% curcuma meal (151.6 mm/g/10sec). The results of the study indicated that there is beneficial effect of dietary inclusion of curcuma (*Curcuma longa*) powder at 0, 2.0, 4.0 and 6.0 per cent on meat fat and meat tenderness of broiler.

Keywords : curcuma meal, meat fat, meat tenderness

Introduction

Texture of foods is mostly determined by the moisture and fat contents, and the types and amounts of structural carbohydrates and proteins. Changes in texture are caused by loss of moisture or fat, and coagulation or hydrolysis of protein (Fellows, 1990). Lean meat contain a very high amount of protein and water and very little fat. In chicken, the protein content are 21 g/100 g; fat 3 g/100g and water 75 g/100g (Simonsen, et al, 1988 in Cross and Overby, 1988). Meat usually cooked before being eaten, and in gastronomic terms, meat is rare if cooked to an internal temperature of 60°C and well done if cooked at 80°C. The endomysial tissue begin to shrink at 50°C and complete at 70°C. When the heating prolonged, the collagen fibres become swollen and gradually denature; and above 70°C will caused further disruption of collagen, and eventually the collagen will solubilized as gelatin. The fat content of meat and meat products has to be judged primarily from the standpoint of calories (Hostetler and Landmann, 1968; Schmidt and Parrish, 1971 in Cross and Overby, 1988). Tenderness is the process of partial relaxation of the fibres. Resolution of rigor is due to enzymatic activity and physical stretching of the muscles fibres attached to bones. Tenderness is measured by use of specialized laboratory equipment or by a taste-panel (Bell and Weaver, 2002). Contrary to popular belief, what the animal is fed does not directly influence tenderness. Many factor influence meat tenderness. The most important factors are genetics, age of the animal, location of the cut on the carcass, processing, method of cooking and degree of doneness (Epley, 2011).

Curcuma (Zingiberaceae) is a large genus of rhizomatousherbs distributed in tropical and subtropical regions especially in India, Thailand, the Malay Archipelago, Indochina, Northern Australia and Indonesia. Many species have been cultivated, and their powdered rhizomes have been widely used as flavours in native dishes and ingredients in many traditional medicines to treat various ailments (Jantan, et al, 2012). *Curcuma longa* has significantly greater total polyphenols, flavonoids and anthocyanidins and anti-oxidant activity (Trinidad et al, 2012). According to Jantan et al, (2012) the three curcuminoids

showed strong inhibition on LDL peroxidation, with curcumin and demethoxycurcumin showing comparable antioxidant activity and more potent than bisdemethoxycurcumin. The three curcuminoids showed strong inhibition on LDL peroxidation. The present study was in accordance with previous studies which indicated that the absence of one methoxy group (demethoxycurcumin) on the phenyl ring did not have effect, but the absence of both methoxy groups (bisdemethoxycurcumin) resulted in decreased antioxidant activity in curcuminoids. The phenolic hydroxyl and the methoxyl groups on the phenyl ring and the 1,3-diketonesystem are important structural features for antioxidant activity. The fermentable dietary fiber from *Zingiberofficinale* and *Curcuma longa* was shown to produce only the short chain fatty acid, propionate which was significant for both samples indicating protective effect for cholesterol-lowering. Propionate release in the colon after dietary fiber fermentation is readily taken up by the liver. Its action is to inhibit the limiting enzyme HMG Co-enzyme reductase for cholesterol synthesis (Trinidad, et al, 2012). According to Basavaraj (2011), results of meat parameters such as live weight (g), and carcass weight (g), dressing percent, meat to bone ratio are lower and chemical composition of meat, there is no significance between control and curcuma treatments, because the broiler rabbit are on summer stress.

Materials and Methods

Forty broilers, day old chicks Arbor Acres CP-707 were assigned randomly and studied for six weeks. Research using Completely Randomized Design (CRD). They were randomly allotted to four dietary treatment groups of ten chicken broilers in each group namely R-0, R-1, R-2 and R-3. The dietary treatments are: R-0 basal diet as control (0% curcuma meal), R-1 basal diet + 2% curcuma meal, R-2 basal diet + 4% curcuma meal and R-3 basal diet + 6% curcuma meal, and each treatment were repeated five times. Carcass composition of meat were analyzed for the fat content. The broiler carcass tenderness was established by meat tenderness instruments.

Results and Discussions

The effect of Curcuma meal on broiler meat fat.

In Table 1, there are the results of the effect of curcuma meal in ration on broiler meat fat.

Table 1. The effect of curcuma meal in ration on broiler meat fat (%)

Replication	R-0	R-1	R-2	R-3
I	2.01	1.92	1.69	1.64
II	1.99	1.93	1.73	1.64
III	1.97	1.91	1.70	1.66
IV	1.98	1.91	1.71	1.65
V	2.00	1.93	1.72	1.66
Average	1.99	1.92	1.71	1.65

Notes :

R-0 basal diet as control – 0% curcuma meal,

R-1 basal diet + 2% curcuma meal,

R-2 basal diet + 4% curcuma meal and

R-3 basal diet + 6% curcuma meal

From Table 1, the average of meat fat are between 1.65% to 1.99%. The highest meat fat is from R-0 the basal diet without curcuma meal (1.99%) and the lowest is from R-3 that using basal diet plus 6% curcuma meal (1.65%). It means that adding curcuma meal in the ration will give less meat fat to the broiler. According to Jantan et al, 2012 the three curcuminoids showed strong inhibition on LDL peroxidation, with curcumin and demethoxycurcumin showing comparable antioxidant activity and more potent than bisdemethoxycurcumin. The fermentable dietary fiber from *Curcuma longa* was shown to produce only the short chain fatty acid, propionate which was significant for the samples indicating protective effect for cholesterol-lowering. Propionate release in the colon after dietary fiber fermentation is readily taken up by the liver. Its action is to inhibit the limiting enzyme HMG Co-enzyme reductase for cholesterol synthesis (Trinidad, et al, 2012).

The effect of Curcuma meal on broiler carcass tenderness.

In Table 2, there are the results from using of curcuma meal in ration, to the broiler carcass tenderness. The highest carcass tenderness was get from the broiler that fed R-3 basal diet with 6% curcuma meal (187.2 mm/g/10sec) and the lowest was get from the broiler that fed basal diet R-0 (151.6 mm/g/10sec).

Table 2. The effect of curcuma meal in ration on broiler carcass tenderness (mm/g/10 sec)

Replication	R-0	R-1	R-2	R-3
I	146.0	141.0	182.0	192.0
II	162.0	171.0	178.0	195.0
III	142.0	154.0	175.0	185.0
IV	149.0	177.0	185.0	182.0
V	159.0	162.0	172.0	184.0
Average	151.6	163.0	178.4	187.2

Notes :

R-0 basal diet + 0% curcuma meal, as control,

R-1 basal diet + 2% curcuma meal,

R-2 basal diet + 4% curcuma meal and

R-3 basal diet + 6% curcuma meal.

From Table 2, adding curcuma meal has effect to the broiler carcass tenderness. And the tenderness will increase when the curcuma meal level percentage more higher. In R-0 (basal diet + 0% curcuma meal), the tenderness are 151.6 mm/g/10sec, will increase when the level of curcuma meal are 2% (R-1 = 163.0 mm/g/10sec); and in R-2 (basal diet + 4% curcuma meal) the tenderness is 178.4 mm/g/10sec; compared to the R-3 basal diet + 6% curcuma meal in ration (187.2 mm/g/10sec). Tenderness of the carcass was increase, because the curcuma has enzymatic activities. Because of the activities of the enzyme, even the fat content in the meat decreased, but the meat tenderness will higher than the meat with no curcuma in the ration. Tenderness is the process of partial relaxation of the fibres. Resolution of rigor is due to enzymatic activity and physical stretching of the muscles fibres attached to bones (Bell and Weaver, 2002). According to Jantan et al, 2012 the three curcuminoids showed strong inhibition on LDL peroxidation, with curcumin and demethoxycurcumin showing comparable antioxidant activity and more potent than bisdemethoxycurcumin.

Conclusions

The effect of curcuma in the ration on poultry meat, will decreased the meat fat, but it also will increased the tenderness of poultry meat. So, the curcuma can be used as natural tenderizer on poultry meat.

References

1. Basavaraj, M., Nagabhushana, V*, Prakash, N., Appannavar, M.M., Prashanth Wagmare and S.Mallikarjunappa. 2011. Effect of Dietary Supplementation of *Curcuma Longa* on the Biochemical Profile and Meat Characteristics of Broiler Rabbits under Summer Stress. *Veterinary World* 2011, Vol.4 (1):15-18
2. Bell D. D., William D.W. Jr.(2002): *Commercial Chicken Meat and Egg Production*. 5th ed. Kluiver Academic Publishers, Massachusetts, 934.
3. Cross, H.R. and A. J. Overby, 1988. *Meat Science, Milk Science and Technology*. Elsevier, Amsterdam. p.23.
4. Epley R.J. (2011): *Meat Tenderness*. Regent of The University of Minnesota.
5. Fellows, P. J. 1990. *Food Processing Technology. Principles and Practice*. Ellis Horwood. New York. p.31
6. Jantan, I, F. C. Saputri, M. N. Qaisar, and F. Buang, 2012. Correlation between Chemical Composition of *Curcuma domestica* and *Curcuma xanthorrhiza* and Their Antioxidant Effect on Human Low-Density Lipoprotein Oxidation. *Hindawi Publishing Corporation Evidence-Based Complementary and Alternative Medicine Volume 2012, Article ID 438356*, doi:10.1155/2012/438356
7. Simonsen, B., R. Hamm, and B. Rogowski. 1988. *Meat as Food*. In. Cross, H.R. and A. J. Overby, 1988. *Meat Science, Milk Science and Technology*. Elsevier, Amsterdam. p.128.
8. Trinidad P. T., R. S. Sagum, M. P. de Leon, A. C. Mallillin, M. P. Borlagdan.2012. *Zingiber Officinale and Curcuma Longa as Potential Functional Foods/Ingredients Food and Public Health* 2012, 2(2): 1-4 DOI: 10.5923/j.fph.20120202.01