



Makara Journal of Science

Editor in Chief

1. [Tribidasari Anggraningrum Ivandini](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Indonesia

Associate Editors

1. [Andi Salamah](#), Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Indonesia
2. [Andreas Martin](#), Leibniz Institute for Catalysis, University of Rostock (LIKAT), Germany
3. [Andrivo Rusydi](#), Department of Physics, Faculty of Science, National University of Singapore, Singapore
4. [Angela Köckritz](#), Leibniz Institut für Catalysis, University of Rostock (LIKAT), Germany
5. [Arben Merkoci](#), Catalan Institute of Nanoscience and Nanotechnology, Spain
6. [Brandon Izak Samuel van der Ventel](#), Department of Physics, Stellenbosch University, South Africa
7. [Gunawan Indrayanto](#), Plant Biotechnology Research Group, Faculty of Pharmacy, Universitas Airlangga, Indonesia
8. [Hans Werner Hammer](#), Institut für Kernphysik, TU Darmstadt, Germany
9. [Iman Hidayat](#), Microbiology Division, Research Center for Biology - LIPI, Indonesia
10. [Iskhaq Iskandar](#), Department of Physics, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Palembang, Indonesia
11. [Jarnuzi Gunlazuardi](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Indonesia
12. [Jeong-Myeong Ha](#), Clean Energy Research Center, Korea Institute of Science and Technology, Korea, Republic Of
13. [Makoto Kakishima](#), Jilin Agricultural University, Engineering Research Center of Chinese Ministry of Education for Edible and Medicinal Fungi, China
14. [Oki Muraza](#), Department of Chemical Engineering, King Fahd University of Petroleum and Minerals, Saudi Arabia
15. [Russell F Howe](#), Department of Chemistry, University of Aberdeen, United Kingdom
16. [Susumu Kajiwara](#), Graduate School of Bioscience and Biotechnology, Tokyo Institute of Technology, Tokyo, Japan
17. [Terry Mart](#), Department of Physics, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Indonesia
18. [Teruna Jaya Siahaan](#), Department of Pharmaceutical Chemistry, The University of Kansas, United States
19. [Wellyzar Sjamsuridzal](#), Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Indonesia
20. [Yuni Krisyuningsih Krisnandi](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Indonesia

Layout Editor

1. [Muklis Sutami](#), Directorate of Research and Community Engagement, Universitas Indonesia, Depok, Indonesia, Indonesia

Vol 21, No 4 (2017): December

Articles

- [High Levels of Resistance in A Culex quinquefasciatus Population to the Insecticide Permethrin in Filariasis Endemic Areas in Central Java](#) [PD](#)
[F](#)
p. 149-154 Irfanul Chakim, Sayono Sayono, Rahayu Astuti 10.7454/mss.v21i4.5664
- [Identification of A Major Quantitative Trait Locus for Grain Weight In Rice Using Microsatellite Marker](#) [PD](#)
[F](#)
p. 155-162 Mariana Susilowati, Hajrial Aswidinnoor, Wening Enggarini, Kurniawan Rudi Trijatmiko 10.7454/mss.v21i4.6590
- [Activity of Cytotoxic Flavanoids against a P-388 Murine Leukemia Cell Line from the Stem Bark of Aglaia elliptica \(Meliaceae\)](#) [PD](#)
[F](#)
p. 163-168 Ace Tatang Hidayat, Kindi Farabi, Desi Harneti, Rani Maharani, Nurlelasari Nurlelasari, Unang Supratman, Yoshihito Shiono 10.7454/mss.v21i4.8836
- [Identification of Avian Influenza Genetic Resistance Gene Marker in Chickens](#) [PD](#)
[F](#)
p. 169-174 R Susanti, Nur Rahayu Utami, Nur Rahayu Utami, Margareta Rahayuningsih, Fidia Fibriana, Margareta Rahayuningsih, Fidia Fibriana 10.7454/mss.v21i4.7428
- [UV-Visible Optical Absorbance of Graphene Oxide Synthesized from Zinc-Carbon Battery Waste via a Custom-Made Ultrasound Generator based on Liquid Sonication Exfoliation Method](#) [PD](#)
[F](#)
p. 175-181 Rhyko Irawan Wisnuwijaya, Agus Purwanto, Wipsar Sunu Brams Dwandaru 10.7454/mss.v21i4.6752
- [Identification of Massoia Lactone and its Hydroxy-derivative from Kabatiella caulivora, an Endophyte of the Terrestrial Plant Alyxia reinwardtii](#) [PD](#)
[F](#)
p. 182-186 Izzatul Laili, Sri Gunarso, Nashrul Wathan, Noor Erma Sugijanto, Gunawan Indrayanto 10.7454/mss.v21i4.8840
- [The Abundance of Nitrogen Fixing, Nitrifying, Denitrifying and Ammonifying Bacteria in the Soil of Tropical Rainforests and Oil Palm Plantations in Jambi](#) [PD](#)
[F](#)
p. 187-193 Zulfarina Zulfarina, Iman Rusmana, Nisa Rachmania Mubarik, Dwi Andreas Santosa 10.7454/mss.v21i4.8841

MAKARA Journal of Science publishes high quality research on fundamental science and provides a broad-based forum for the information sharing of ongoing research and development in science. Topics covered include physics, chemistry and biology. All submitted manuscripts will be subjected to the peer review process following a thorough inspection by the editors. Makara Journal of Science is the successor to the "MAKARA of Science Series" and "Makara Sains", which have been published since 1997. Starting from 2014, the journal has been periodically published four times a year (March, June, September, and December). Publication in this journal is free of charge and all published papers are available online through open access.

[Read more](#)

Editor in Chief [Tribidasari Anggraningrum Ivandini](#) | [Full editorial board](#)

[Statistics](#) | [Citations](#)

catechin (**1a**), epicatechin (**1b**), gallicocatechin (**2a**), and epigallocatechin (**2b**), together with their cytotoxic activity against murine leukemia cells.

Materials and Methods

Equipment. UV spectra were measured using a Shimadzu UV-160A ultraviolet-visible spectrometer, with MeOH (Kyoto, Japan). The IR spectra were recorded on a Perkin-Elmer 1760X FT-IR in KBr (Waltham, MA, USA). The mass spectra were recorded with a Synapt G2 mass spectrometer instrument (Waters, Milford, MA, USA). NMR data were recorded on a JEOL ECZ-600 spectrometer at 600 MHz for ^1H and 150 MHz for ^{13}C (Tokyo, Japan), using TMS as an internal standard. Column chromatography was conducted on silica gel 60 (Kanto Chemical Co., Inc., Japan). TLC plates were pre-coated with silica gel GF₂₅₄ (Merck, 0.25 mm), and detection was achieved by spraying with 10% H₂SO₄ in EtOH, followed by heating.

Plant material. The stem bark of *A. elliptica* was collected in Bogor Botanical Garden, Bogor, West Java Province, Indonesia, in June 2015. The plant was identified by the staff of the Bogoriense Herbarium, Bogor, and a voucher specimen (No. Bo-1294562) was deposited at the Herbarium.

Plant extraction. Dried ground stem bark (2.3 kg) of *A. elliptica* was extracted with methanol (12 L) at room temperature for 3 days. After removal of the solvent under vacuum, the viscous concentrate of MeOH extract (321.5 g) was first suspended in H₂O and then partitioned successively with *n*-hexane, EtOAc, and *n*-butanol. Evaporation resulted in the crude extracts of *n*-hexane (22.6 g), EtOAc (31.4 g), and *n*-butanol (34.5 g), respectively. The *n*-hexane, EtOAc, and *n*-butanol extracts exhibited cytotoxic activity against P-388 murine leukemia cells, with IC₅₀ values of 67.72, 32.69, and >100 µg/mL, respectively. The EtOAc soluble fraction (20 g) was fractionated by column chromatography on silica gel using a gradient *n*-hexane and EtOAc to give fractions A–E, combined according to the TLC results. Fraction D (1.73 g) was subjected to column chromatography over silica gel, using a gradient mixture of CHCl₃:Me₂CO (10:0–1:1) as eluting solvents to afford six subfractions (D1–D6). Subfraction D4 (460 mg) was chromatographed on a column of silica gel, eluted with CHCl₃–MeOH (10:0–4:1), to give five subfractions (D4A–D4E). Subfraction D4D was chromatographed on preparative TLC, eluted with CHCl₃–MeOH (8.5:1.5), to give compound **1** (47.5 mg). Subfraction D5 (600 mg) was chromatographed on a column of silica gel, eluted with CHCl₃–MeOH (10:0–7:3), to give five subfractions (D5A–D5E). Subfraction D5D was chromatographed on a column of silica gel, eluted with CHCl₃–MeOH (10:0–1:1), to give compound **2** (127 mg).

Determination of cytotoxic activities [8,11,20]. The P-388 cells were seeded into 96-well plates at an initial cell density of approximately 3×10^4 cells cm⁻³. After 24 h of incubation for cell attachment and growth, varying concentrations of samples were added. The compounds added were first dissolved in DMSO at the required concentration. Six subsequent desirable concentrations were prepared using phosphoric buffer solution (PBS, pH=7.30–7.65). Control wells received only DMSO. The assay was terminated after a 48 h incubation period by adding MTT reagent (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide, also named thiazol blue), and the incubation was continued for another 4 h, in which the MTT-stop solution containing sodium dodecyl sulphate (SDS) was added, and another 24 h incubation was conducted. Optical density was read using a micro plate reader at 550 nm. IC₅₀ values were taken from the plotted graph of the percentage of live cells compared to control (%) receiving only PBS and DMSO, versus the tested concentration of compounds (µM). The IC₅₀ value is the concentration required for 50% growth inhibition. Each assay and analysis was run in triplicate and averaged.

Results and Discussion

The stem bark of *A. elliptica* was ground and successively extracted with MeOH and partitioned with *n*-hexane, ethyl acetate, and *n*-butanol. All the extracts were evaluated for cytotoxic activity against P-388 murine leukemia cells. The ethyl acetate extracts exhibited the strongest cytotoxic activity against P-388 murine leukemia cells, with IC₅₀ values of 32.69 µg/mL. Subsequent phytochemical analysis was therefore focused on the EtOAc extract of *A. elliptica*. The EtOAc extract was chromatographed over a column packed with silica gel 60 by gradient elution. The fractions were repeatedly subjected to normal-phase column chromatography to afford flavonoid compounds **1** and **2** (Figure 1).

Mixture of catechin and epicatechin (1), yellow amorphous powder, m.p. 176–177 °C, UV (MeOH) λ_{max} nm (log ε) 275 (3.93), IR (KBr) ν_{max}(cm⁻¹) 3330 (O-H stretch), 1572 (C=C ring stretch), 1146 (asymmetric C-O-C stretch), 1051 (symmetric C-O-C stretch), 827 (substituted benzene ring). $^1\text{H-NMR}$ (CD₃OD, 600 MHz), see Table 1; $^{13}\text{C-NMR}$ (CD₃OD, 125 MHz), see Table 1; HR-TOFMS (positive ion mode) *m/z* 291.0878 [M+H]⁺ (calcd. for C₁₅H₁₄O₆, *m/z* 290.0790).

Mixture of gallicocatechin and epigallocatechin (2), brown amorphous powder, m.p. 198–202 °C, UV (MeOH) λ_{max} nm (log ε) 277 (4.07), IR (KBr) ν_{max}(cm⁻¹) 3325 (O-H stretch), 1577 (C=C ring stretch), 1144 (asymmetric C-O-C stretch), 1062 (symmetric C-O-C stretch), 829 (substituted benzene ring). $^1\text{H-NMR}$ (CD₃OD, 600 MHz), see Table 1; $^{13}\text{C-NMR}$ (CD₃OD,

Activity of Cytotoxic Flavanoids against a P-388 Murine Leukemia Cell Line from the Stem Bark of *Aglaia elliptica* (Meliaceae)

Ace Tatang Hidayat^{1,2}, Kindi Farabi¹, Desi Harneti¹, Rani Maharani^{1,2}, Nurlelasari¹, Unang Supratman^{1,2*}, and Yoshihito Shiono³

1. Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Jatinangor 45363, Indonesia

2. Central Laboratory of Universitas Padjadjaran, Jatinangor 45363, Indonesia

3. Department of Food, Life, and Environmental Science, Faculty of Agriculture, Yamagata University, Tsuruoka, Yamagata 997-8555, Japan

*E-mail: unang.supratman@unpad.ac.id

Received May 16, 2017 | Accepted November 9, 2017

Abstract

Two mixtures of flavanoid compounds (**1** and **2**), a mixture of catechin (**1a**) and epicatechin (**1b**), and a mixture of gallo catechin (**2a**) and epigallo catechin (**2b**), were isolated from the active fraction of the stem bark of *Aglaia elliptica* methanol extract. The chemical structure of the compounds was identified with spectroscopic data, including UV, IR, NMR (¹H, ¹³C, DEPT 135°, HMQC, HMBC, ¹H-¹H COSY), and MS, and additionally compared with previously reported spectral data. All compounds were evaluated for their cytotoxic effects against P-388 murine leukemia cells. Compound **2** showed cytotoxicity against the P-388 murine leukemia cell, with an IC₅₀ value of 7.79 µg/mL, but compound **1** was found not to be active (more than 100 µg/mL).

Abstrak

Aktivitas Sitotoksik Senyawa Flavanoid Terhadap Sel Murine Leukemia P-388 dari Kulit Batang *Aglaia elliptica* (Meliaceae). Dua campuran senyawa flavanoid (1 dan 2), suatu campuran dari katekin (1a) dan epikatekin (1b) serta campuran dari gallo katekin (2a) dan epigallo katekin (2b) telah diisolasi dari kulit batang *Aglaia elliptica*. Struktur kimia senyawa tersebut diidentifikasi berdasarkan data spektroskopi, meliputi UV, IR, NMR (¹H, ¹³C, DEPT 135°, HMQC, HMBC, ¹H-¹H COSY) dan MS, serta tambahan dengan perbandingan data spektra yang diperoleh sebelumnya. Semua senyawa dievaluasi aktivitas sitotoksiknya terhadap sel murin leukemia P-388. Senyawa 2 menunjukkan aktivitas sitotoksik terhadap sel murin leukemia P-388 dengan nilai IC₅₀ 7,79 µg/mL, sedangkan senyawa 1 tidak memberikan aktivitas sitotoksik (IC₅₀ lebih besar dari 100 µg/mL).

Keywords: *Aglaia elliptica*, cytotoxic activity, flavanoid, Meliaceae, sel murine leukemia P-388

Introduction

Aglaia is distributed mainly in tropical rainforests of the Indo-Malaysian region [1]. The genus *Aglaia* (Meliaceae) is the largest genus of the Meliaceae family, comprising more than 150 species, approximately 65 of which grow in Indonesia [1,2]. Extracts from the *Aglaia* genus have been used traditionally for treating certain diseases. In Thailand, *Aglaia odorata* is used to treat heart disease, bruises, traumatic injury, febrifuge, and toxins, by causing vomiting [3]. Previous phytochemical studies of this genus revealed the presence of a compound with interesting biological activity, including antifungal and

antitumor sesquiterpenoids [4,5], cytotoxic and anti-inflammatory diterpenoids [3,6], cytotoxic and anti-retroviral triterpenoids [7-10], cytotoxic steroids [4], cytotoxic and anti-inflammatory alkaloids [10,11,14], and cytotoxic rogamides [13,14].

A. elliptica is a tree, mainly distributed in the northern part of Sulawesi in Indonesia [15]. Previous phytochemical study of this plant reported the presence of diamide and cycloartane-type triterpenoid from the leaves [17] and novel cytotoxic 1*H*-cyclopenta[*b*]benzo furan from the fruits [16]. In the present paper, we elucidate the isolation and structure of the mixture of flavonoid compounds,