


MAF 12

12th International Conference on Methods and Applications of Fluorescence

SPECTROSCOPY, IMAGING AND PROBES

Strasbourg, France
September 11-14, 2011

BOOK OF ABSTRACTS



COMMITTEES

International Program Committee

| | |
|-------------------------------------|---------------------------------|
| Otto S. Wolfbeis (Germany) | Yun-Bao Jiang (China) |
| A. Ulises Acuña (Spain) | Lennart B.Å. Johansson (Sweden) |
| Manfred Auer (UK) | Anita C. Jones (UK) |
| Mario Berberan-Santos (Portugal) | Paavo K.J. Kinnunen (Finland) |
| David J.S. Birch (UK) | Joseph R. Lakowicz (USA) |
| John Birmingham (USA) | Helge Lemmetyinen (Finland) |
| Amilra P. de Silva (UK) | János Matkó (Hungary) |
| Alexander P. Demchenko (Ukraine) | Yves Mély (France) |
| Alberto Diaspro (Italy) | Ute Resch-Genger (Germany) |
| Jörg Enderlein (Germany) | Catherine Royer (France) |
| Hans C. Gerritsen (The Netherlands) | Petra Schwillle (Germany) |
| Enrico Gratton (USA) | Claus Seidel (Germany) |
| Albin Hermetter (Austria) | Trevor Smith (Australia) |
| Martin Hof (Czech Republic) | Hans J. Tanke (The Netherlands) |
| Johan Hofkens (Belgium) | Jacek Waluk (Poland) |
| Totaro Imasaka (Japan) | Jerker Widengren (Sweden) |
| David M. Jameson (USA) | Sergiy M. Yarmoluk (Ukraine) |

Local Organizing Committee

Yves Mély (Chair)
Guy Duportail (Co-chair)
Cyril Favard
Andrey Klymchenko
Youri Arntz
Frédéric Bolze
Pascal Didier
Marlyse Wernert (secretary)
Ingrid Barthel (secretary)
Ludovic Four (webmaster)

- P149** *Fluorescence imaging of (phospho)lipase patterns using differential activity-based gel electrophoresis (DABGE)*
 Maria Morak, Bojana Stojić, Albin Hermetter
- P150** *Conformational transitions in the Mg²⁺-dependent 7S11 deoxyribozyme, a fluorescence study*
Elisa Turriani, Claudia Höbartner, Thomas M. Jovin
- P151** *The heart tissues differentiation by time resolved fluorescence spectroscopy*
Jonas Venius, Edvardas Žurauskas, Ričardas Rotomskis
- P152** *Organization of ECF and ABC Transporter Modules During Substrate Translocation*
Joanna Ziolkowska, Johanna Heuveling, Daniela Weidlich, Franziska Kirsch, Thomas Eitinger, Erwin Schneider, Andreas Herrmann
- P153** *The influence of polymer structure on the cationic polymer – DNA complexes revealed by time-resolved fluorescence studies*
Tiia-Maaria Ketola, Martina Hanzlíková, Linda Cameron, Arto Urtti, Helge Lemmetyinen, Marjo Yliperttula, Elna Vuorimaa
- P154** *Poly(β -amino ester) - DNA complexes: time-resolved fluorescence and cellular transfection studies*
Elna Vuorimaa-Laukkanen, Tiia-Maaria Ketola, Jordan J. Green, Martina Hanzlíková, Helge Lemmetyinen, Robert Langer, Daniel G. Anderson, Arto Urtti, Marjo Yliperttula
- P155** *Modulation of raft-dependent immune functions by a monoclonal cholesterol-specific antibody: a biophysical study on the mechanism of action*
Andrea Balogh, Emese Izsépi, Anikó Molnár, László Cervenák, George Füst, Zoltán Beck, György Vámosi, Gábor Mocsár, Glória László, János Matk
- P156** *The action of chlorogenic acid on erythrocyte and model membranes*
Dorota Bonarska-Kujawa, Hanna Pruchnik, Halina Kleszczyńska
- P157** *Fluorescence studies on new potential antitumoral 1,3-diarylurea derivatives of thieno[3,2-b]pyridines in solution and in nanoliposomes*
Elisabete M. S. Castanheira, Cátia N. C. Costa, Ana Rita O. Rodrigues, Daniela Peixoto, Maria-João R. P. Queiroz
- P158** *Study of morphology, hydration and mobility of cationic lipophosphoramidates liposomes by fluorescent solvent relaxation*
 Damien Loizeau, Piotr Jurkiewicz, Laure Deschamps, Paul-Alain Jaffrès, Martin Hof, Philippe Giamarchi
- P159** *Characterization by steady-state and time resolved fluorescence spectroscopy of the lipid and protein organization in HBsAg particles*
Vanille J. Greiner, Caroline Egelé, Sule Oncul, Frédéric Ronzon, Catherine Manin, Andrey Klymchenko, Yves Mély
- P160** *Spectroscopic studies of the novel, cationic gluconamide-type surfactants/DNA and lipid interactions*
Teresa Kral, Hanna Pruchnik, Martin Hof, Renata Skrzela, Kazimiera A. Wilk
- P161** *Effects of supplementation with L-proline or inositol on yeast membrane fluidity and ethanol tolerance*
 Safri Ishmayana, Ursula Kennedy, Robert Learmonth
- P162** *Study of the phase transition of mixtures of original cationic phospholipids by fluorescence anisotropy*
Sélim Mahfoudhi, Damien Loizeau, Laure Deschamps, Paul-Alain Jaffrès, Tristan Montier, Pierre Lehn, Philippe Giamarchi
- P163** *Dynamic organization of HIV co-receptors in the membrane of immune cells revealed by single quantum dot tracking*
Patrice Mascalchi, Anne-Sophie Lamort, Evert Haanapel, Fabrice Dumas, Bernard Lagane, Fernando Arenzana-Seisdedos, André Lopez, Laurence Salomé
- P164** *Lipid raft detecting in membranes of live erythrocytes*
Ilya Mikhalyov, Andrey Samsonov

Effects of supplementation with L-proline or inositol on yeast membrane fluidity and ethanol tolerance

Safri Ishmayana^{1,2}, Ursula Kennedy¹ & Robert Learmonth^{1*}

¹Centre for Systems Biology, University of Southern Queensland, Toowoomba 4350 (Australia)

²Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Bandung (Indonesia)

The decrease of fossil fuel availability has created a high demand for alternative fuels, including bioethanol produced by yeast fermentation of carbohydrate. Relatively low ethanol yields can be problematic, and approaches to increase efficiency have included genetic modification of yeast to improve metabolic flux, fermentation rate and ethanol tolerance. We investigated an alternative approach, aiming to evaluate novel yeasts for bioethanol production, and further to enhance their ethanol tolerance through modification of growth medium composition. We focused specifically on two important components which have been reported to positively affect yeast stress tolerance; the sugar inositol^[1,2] and the amino acid L-proline^[3]. Three yeast strains (a bakers' yeast, a wine yeast and a sake yeast), known tolerate 17% ethanol, were studied under aerobic conditions. Cultures were supplemented with various concentrations of either inositol or L-proline, and sampled during respiro-fermentative growth (fermentation of glucose to ethanol) and respiratory growth on ethanol. Physiological parameters, viability, membrane fluidity (by laurdan GP) and tolerance to 18% ethanol were determined. Differences were noted in the growth, ethanol productivity and ethanol tolerance of the yeasts, with significant differences evident in laurdan GP in respiro-fermentative, but not respiratory phase cells. As expected, tolerance to ethanol was significantly higher in respiratory phase cells, also correlating with significantly higher laurdan GP which indicated lower membrane fluidity. However, on trialling L-proline levels from 0.1 to 3 g/L and inositol levels from 0.002 to 0.2 g/L, no significant differences were found in laurdan GP or ethanol tolerance. Thus we could not confirm previous reports of positive effects of L-proline or inositol, due largely to high variability in the data from triplicate experiments. It is possible that the effects of these supplements may be strain dependant; the wine yeast tended towards lower membrane fluidity with 0.5 g/L proline, and in the sake yeast there was a trend towards improved ethanol tolerance with inositol levels above 0.005 g/L. Future experiments will investigate a wider variety of yeast strains and increase repetition to minimise data variability.

The authors wish to acknowledge the support of a scholarship from the Ministry of National Education of the Republic of Indonesia to enable postgraduate study by Safri Ishmayana.

References: [1] E. L. Krause et al., *Indust. Biotechnol.* 3 (2007) 260. [2] R. Ji et al., *J. Agro-Environ. Sci.* 27 (2008) 2080. [3] H. Takagi et al., *Appl. Env. Microbiol.* 71 (2005) 8656.

*Corresponding author: e-mail: Robert.Learmonth@usq.edu.au